

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**


Defective images within this document are accurate representations of the original documents submitted by the applicant.


Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**





Eur päisches Pat ntamt  
Europ an Pat nt Offic  
Offic europé n d s br v ts



(11) **EP 1 038 959 A1**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:  
27.09.2000 Bulletin 2000/39

(51) Int Cl.7: **C12N 15/12, C07K 14/755**

(21) Application number: **99104050.2**

(22) Date of filing: **17.03.1999**

(84) Designated Contracting States:  
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU  
MC NL PT SE**  
Designated Extension States:  
**AL LT LV MK RO SI**

(72) Inventors:  
• **Négrier, Claude, Dr.**  
**69540 Irigny (FR)**  
• **Plantier, Jean Luc**  
**69520 Grigny (FR)**

(71) Applicant: **Aventis Behring Gesellschaft mit  
beschränkter Haftung**  
**35002 Marburg (DE)**

(54) **Factor VIII without B-domain, comprising one or more insertions of a truncated intron I of factor IX**

(57) A modified Factor VIII cDNA is described in which the B-domain of the wild-type Factor VIII cDNA has been deleted and a truncated Factor IX intron 1 has

been inserted in one or more locatons of the Factor VIII cDNA. Such modified Factor VIII cDNA may be used for a higher yield production of Factor VIII in vitro as well as in a transvector for gene therapy.

EP 1 038 959 A1

## Description

[0001] This invention is directed to a modified Factor VIII cDNA and its use for the improvement of the Factor VIII production.

5 [0002] Factor VIII (FVIII) is a plasma coagulation cofactor implicated in the activation of Factor X (FX). A decrease in the presence or activity of Factor VIII in blood stream leads to hemophilia A. The level of the decrease in Factor VIII activity is directly proportional to the severity of the disease (Foster and T.S., 1989; Kaufman, 1992; Vlot et al., 1998). The current treatment of hemophilia A consists of the replacement of the missing protein by plasma-derived or recombinant FVIII. Recombinant FVIII is produced in CHO or BHK cells after selection of the best producing clones and amplification of the FVIII cDNA copy number.

10 [0003] Several studies have stressed the low FVIII production level in different cellular systems. Biosynthesis of FVIII was shown to be regulated in at least three different levels. First, among the FVIII cDNA sequence two nucleotides stretches, localized in the A2 coding domain, were demonstrated to act as transcriptional silencers (Fallaux et al., 1996; Hoebe et al., 1995; Koeberl et al., 1995; Lynch et al., 1993). Second, FVIII protein synthesis is tightly regulated by several reticulum endoplasmic chaperones (BiP; Calreticulin; Calnexin; ERGIC-53). Many of these interactions retain FVIII in the cell and direct it through the cellular degradation machinery (Dorner et al., 1987; Nichols et al., 1998; Pipe et al., 1998). Third, once secreted FVIII is sensitive to protease degradation and needs to be protected by von Willebrand Factor (vWF) (Kaufman et al., 1989).

20 [0004] It is therefore a problem to develop improved processes which result in higher yields of FVIII. The present invention offers a solution to this problem by a modified FVIII cDNA.

[0005] According to this invention a modified FVIII cDNA is made available in which the B-domain of the wild-type FVIII cDNA has been deleted and a truncated FIX intron 1 has been inserted in one or more locations of the FVIII cDNA. In addition the B-domain of the wild-type FVIII cDNA has been replaced by four arginines.

[0006] These FVIII constructs were prepared as follows:

#### 1. FVIII cloning

30 [0007] A PCR cloning strategy was designed, based on the synthesis of four PCR fragments, spanning the FVIII cDNA and excepting the B-domain. Based on published data, it was of interest to replace the B-domain by four arginines. Using the MoMuLV reverse transcriptase (Promega, Charbonnières, France) a reverse transcription was done on human cell RNA isolated from a liver biopsy with the written informed consent of the patient. Four PCR fragments were generated using the Expand System (Boehringer-Mannheim, Germany), as described in Table 1.

Table 1:

Summary of FVIII PCR fragments					
Fragment	Oligonucleotide sens antisens	SEQ ID No.	Size in bp	5' cloning site	3' cloning site
1	FVIII ATG 3'Bgl II	No.1 No.2	1704	Not I	Bgl II
2	5'Bgl II 4R AS	No.3 No.4	624	Bgl II	Sal I
3	4R S 3'Bgl I	No.5 No.6	1093	Sal I	Bgl I
4	5'Bgl I FVIII stop	No.7 No.8	1026	Bgl I	Xho I

[0008] All the oligonucleotide sequences used for cloning are shown in Annex 1.

50 [0009] An additional improvement was performed by optimisation of the ATG environment following the rules of Kozak (Kozak, 1997). For this purpose the oligonucleotide FVIII ATG (SEQ ID No 1) was used.

#### Comparison of the wild-type FVIII ATG and the Kozak modified FVIII ATG sequences

55 [0010] WT sequence: TAA GTC ATG CAA ATA

Kozak modified: ACA CCC ATG GAA ATA

[0011] The modified amino-acids are represented in bold.

[0012] Four arginines replace according to the invention the B-domain of the FVIII protein. They are introduced by

the oligonucleotides used for the cloning of the two fragments surrounding the B-domain (see Fragments 2 and 3 of Table I), namely the oligonucleotides 4R AS (SEQ.ID.No.4) and 4R S (SEQ.ID.No.5). The Sal I site was generated by the coding sequence of the arginines as follows:

## SAL I SITE

4R S :5'-A AGA CGT CGA CGA GAA ATA ACT CGT ACT ACT CTT  
4R AS TTG TTA CGG TAA CTT GGT TCT TCT GCA GCT GCT CTT

CORRESPONDING PEPTIDIC SEQUENCE:

Pro Arg Arg Arg Arg Glu Ile Thr Arg Thr Thr Leu

[0013] In the wild-type FVIII the peptidic sequence is:

Pro-Arg-Domain B-Arg-Glu

[0014] This indicates that a Sal I restriction site was inserted through the middle of the fourth arginine coding sequence without any sequence alteration. All PCR fragments were cloned in a pCR2 vector using the T/A cloning kit (InVitrogen, the Netherlands). Two clones of each fragment were entirely sequenced. Some mutations were found in a ratio of 1 for 800 bases. One mutation was silent but three others modified the coding sequence and the three cDNA pieces bearing the mutations were successively exchanged.

[0015] A subsequent sub-cloning strategy leads to the production of two fragments (each being the sum of two PCR products): a 5' 2.3kb (FVIII ATG) and a 3' 2.1kb (FVIII Stop). The 5' 2.3kb (FVIII ATG) and a 3' 2.1kb (FVIII Stop) were constructed and inserted in the expression vector pCDNA3 (InVitrogen, The Netherlands) opened by Not I and Xho I and treated with alkaline phosphatase. The cloning cassette of pCDNA3 presents a similar restriction sequence as pCR2 and the vector possesses its own resistance gene (neomycin). A B-domain deleted FVIII coding cDNA (hereafter referred to Factor VIII cDNA) was directly obtained in the expression vector. The final FVIII cDNA integrity was checked by extensive restriction analysis.

[0016] Factor VIII cDNA was subcloned in the Bluescript pKS II+vector opened by Not I and Xho I. The use of this vector was more convenient for the introduction of subsequent modifications in Factor VIII cDNA (e.g. introns addition; see hereafter).

## 2. Truncated FIX intron 1 insertion

[0017] According to the invention the FVIII cDNA was further modified by the insertion of a Factor IX truncated intron 1 (FIX TI1=SEQ.ID.No.9). The FIXTI1 was inserted in different locations of the FVIII cDNA as follows:

- in the FVIII intron 1 location, to use a similar strategy as for FIX (Kurachi et al., 1995)
- in the FVIII intron 11 and 12 locations, because a transcriptional silencer sequence was described in this region.
- in the FVIII introns 1+11 and 1+12 locations. Since FVIII cDNA is much larger than FIX (4.4 kb vs 1.4 kb), we hypothesized that it might be of interest to introduce supposedly stabilizing sequences at the locations normally occupied by introns 11 and 12 in addition to intron 1. Since the location of introns 11 and 12 are grossly in the middle of the FVIII sequence it is possible that they may act synergistically with intron 1.

[0018] The FIXT11-Sequence (=SEQ-ID No. 9) used according to the invention in different locations of the FVIII cDNA starts after the coding sequence by the splice donor sequence and ends by the splice acceptor sequence of the truncated intron 1. The upper case letters start after and stop before the Nsi I and Mlu I restriction sites, respectively. For details see Annex 2.

5 [0019] For cloning this fragment in an exogenous sequence, two new FVIII junction fragments were generated, one upstream the FIX T11 with the addition of a Nsi I site and one downstream with the addition of a Mlu I site. The three fragments were subsequently linked together using these sites.

[0020] A similar strategy was used for inserting the three FIXT11 in different locations. In each case three PCR fragments (A, B, C) were generated with the Expand System using Factor VIII cDNA as template for segments A and C, and Factor IX intron 1 for the B fragment. The A fragment extremities comprise Factor VIII sequence on the 5' end, and on the 3' end a fusion between the FVIII 3' splicing sequence and the Factor IX first intron 5' splicing sequence. A Nsi I restriction site was added between these two sequences. The B fragment possesses at the 5' extremity a complementary sequence to the previous fragment, the truncated Factor IX intron 1, and at the 3' end an inserted MluI restriction site. The C fragment was made of the complementary sequence of the 3' extremity from fragment B followed by the Factor IX first intron 3' splicing sequence and by the Factor VIII cDNA downstream coding sequence (see Figure 1).

[0021] The B fragment was the same for all constructs whereas A and C were all different, corresponding to the 5' and 3' FVIII sequences of the intron insertion sites.

[0022] The insertion strategies for intron 11 and 12 positions are indicated in Figure 2.

20 [0023] Each PCR fragment was generated and first cloned in pCR2 vector using T/A cloning kit (Invitrogen, The Netherlands). They were linked together in two successive steps (pCR2-B+C/MluI+XhoI; pCR2-A+BC/NsiI+XbaI). All ABC (1,11 and 12) fragments were sequenced and shown to be free of mutations.

Table II:

Summary of the PCR fragments required for introducing TFXI1.	
Fragment Name	Size in bp
I1-A	204
TFXI1 (B)	281
I1-C	464
I11-A	234
I11-C	388
I12-A	434
I12-C	180

40 [0024] ABC1 was cloned in pKS II+ FVIII after Spe I-Spe I digestion. After checking the orientation, the resulting FVIII 11 cDNA was subcloned in pCDNA3 vector using a Not I-Xho I digestion. ABC11 and ABC12 were digested by Bgl II and Sal I and ligated in pKSII-FVIII digested by the same enzymes. Resulting FVIII 111 and FVIII 112 were subcloned in pCDNA3 using a Not I-Xho I restriction.

45 [0025] To generate a construct containing two introns, pKS FVIII 11 and the fragments ABC 11 or ABC 12 were digested by Bgl II and Sal I and ligated together. pKSII FVIII 11+11 and FVIII 11-12 were obtained and FVIII cDNAs containing the introns were subcloned in pCDNA3 by the same strategy as previously described.

### 3. Generation of stably expressing FVIII cell lines

50 [0026] All pCDNA3-FVIII constructs were transfected in CHO cells by electroporation. Briefly,  $7 \times 10^6$  washed CHO cells were electroporated in the presence of 10  $\mu$ g of Pvu I linearized construct. The cells were selected 15 hours after electroporation in IMDM (=Iscove's Modified Dulbecco's Medium) supplemented by antibiotics and 10% fetal calf serum containing 0.6 mg/ml G418. For each construct (FVIII; I1, I11, I12; I1+11; I1+12), two pools of clones containing more than 50 individual colonies were grown and frozen. A set of 25 individual colonies were picked up, grown and assessed for Factor VIII expression using an ELISA method (Asserachrom VIII.C.Ag, Diagnostica STAGO, Asnières, France).

55 For all clones, the transfection efficiency was around 40-60% as determined by the number of expressing clones. The five best FVIII producer clones were kept and frozen. The FVIII antigen detected varied dramatically from one construct to another.

[0027] The activity of the recombinant Factor VIII was measured using the classical one stage clotting assay for each

pool and five selected clones. A coagulant activity was found in each supernatant of Factor VIII expressing cells indicating that the construct was coding for a functional protein. The activity was correlated to the amount of protein detected by ELISA. Therefore, it was demonstrated that each construct allowed the production of a functional procoagulant FVIII.

#### 5 4. Evaluation of the construct efficiency

##### 4.1 Quantitative analysis in CHO cells

[0028] The kinetic of FVIII production was analyzed for each pool of transfected CHO cells. For each pool, independent experiments were made by two different experimentators. At day 0,  $4 \cdot 10^5$  cells were seeded in a 6-well plate in 2 ml of fresh medium containing 0.6 mg/ml G418. From day 1 to day 4, the culture supernatant was collected, centrifuged and assayed for FVIII antigen and procoagulant activity. Each day, after medium removal, cells were trypsinized and counted in Trypan Blue. FVIII accumulation is shown in Figure 3. Three constructs allowed a better production than the non-modified FVIII cDNA, i.e. FVIII I1, FVIII I1+11 and particularly FVIII I1+12. FVIII I1 and FVIII I1+11 led to a 2 to 3 times higher production and FVIII I1+12 to 8 to 9 times more. Procoagulant activities were measured from the same supernatants and are presented in Figure 4. They were directly correlated to the amount of antigen detected.

##### 4.2 Quantitative analysis in HepG2 cells

[0029] To confirm the data obtained in CHO cells, a second cellular model was chosen. The constructs exhibiting an improvement in the FVIII expression were tested (e.g. FVIII; FVIII I1, FVIII I1+11 ; FVIII I1+12) using transiently transfected hepatic cell line HepG2. 2 µg of circular DNA were added to 12 µl of Fugene 6 transfection reagent (Boehringer Mannheim, Meylan, France). After 15 minutes, the complex was added dropwise on a 90 mm dish seeded with  $5 \cdot 10^6$  cells the day before. The transfection mix was incubated for 6h before being replaced by 4 ml of fresh medium. Culture medium was collected 72h later and submitted to coagulation and ELISA assays. As shown in Figure 5, the results obtained using HepG2 cells confirmed the data obtained in CHO cells with the exception of FVIII I1+11 construct which here did not product any detectable FVIII antigen. These data reinforced the potential interest in the FVIII I1+12 construct.

#### 30 5. Analysis of the expression levels

[0030] Total RNA of each transfected pool A was extracted with the Rneasy mini Kit (Quiagen),  $6 \cdot 10^6$  cells being used for each extraction. 10 µg of total RNA were migrated at 120V, 4°C in a 0.8% agarose gel in phosphate buffer, transferred overnight on Hybond-N Nylon membrane (Amersham) and baked for 2h at 80°C. A FVIII RNA probe, containing the 1.03 kb Sal I-Bgl I fragment, was generated by T7 polymerase using the RNA DIG labelling kit (Boehringer). The membrane was blocked 30 min by the Dig easy Hyb solution (Boehringer) and incubated in the same fresh solution overnight with 500 ng of labeled antisense probe. Washes were conducted as recommended by the manufacturer and the blot was revealed using the DIG detection Kit (Boehringer) analysed and quantified. FVIII signals were compared to ribosomal BET-labeled RNA and to GAPDH signal.

[0031] FVIII, FVIII I1 and FVIII I1+11 mRNA amounts were very close to each other. FVIII I1+12 mRNA was expressed in larger amounts. These results indicate a correlation between the amount of mRNA and the protein produced for the constructs (see Figure 6).

[0032] A two time increase in FVIII I1+12 mRNA lead to a 8 to 9 times protein increase. Therefore, the addition of truncated FIX intron I might play a double role in stabilizing FVIII mRNA but also probably in acting during translation.

[0033] The mRNA from FVIII is 4.4 kb long and the differences of 0.3 kb due to a possible non splicing of the TFI1 may not be visible in Northern blot. A RT-PCR set of experiments was done on total RNA extracted from the different cell lines. In each case a band corresponding to a spliced mRNA was obtained indicating the splicing of the TFI1 from FVIII mRNA.

#### 50 6. Protein characterization

[0034] A sheep anti-FVIII antibody from Cedarlane (Hornby, Canada) was purchased and positively tested in a control immunoblot using recombinant FVIII. This antibody was used in immunoblot on cell supernatant but no signal was obtained due to the low amount of secreted antigen. An immunoprecipitation was done on cell supernatant but here again no signal was obtained indicating the inability of this antibody to immunoprecipitate FVIII. An immunoblot was done on Triton-X100 soluble cell lysates. 90mm dishes was lysed with 300 µl ice-cold lysis buffer (Hepes 20mM pH7.5, KC1 100mM, MgCl<sub>2</sub> 2mM, Triton X-100 0.5%). Cells were scrapped and centrifuged at 4°C, 10 min at 14000g. Protein concentration was measured with the Dc-protein Assay kit (BioRad, Hercules, USA). 175 µg of each cell lysates were

loaded on 7.5% acrylamide gel and treated following the Laemmli protocol. After semi-dry transfer (35 min at 400mA), the nitrocellulose membrane was incubated overnight in TBS-T (20mM Tris pH 7.5, NaCl 0.15 M, Tween-20 0.5%). The membrane was then incubated 1h with the anti-FVIII antibody (5 µg/ml) in TBS-T. After three washes of 10 min each in TBS-T, the membrane was incubated for 30 min with a rabbit anti-sheep peroxidase coupled antibody (dilution 10<sup>-4</sup> in TBS-T). Extensive washes were conducted before revelation with the ECL system (Amersham).

[0035] Among all the lysates, only the cells transfected with the FVIII I1+12 construct gave a positive signal with a correct approximative molecular weight. The FVIII I1+12 appeared as a single band product migrating at a molecular weight corresponding to the Factor VIII heavy chain devoided of the B-domain.

[0036] In order to confirm the differences observed in intracellular FVIII amounts, an ELISA was done on Triton-X 100 soluble lysates. The presence of Triton X-100 was shown not to influence the FVIII ELISA test. The value of FVIII antigen present inside the cells confirms the data obtained in Western blot. FVIII I1+12 led to a 100 times higher synthesis of the antigen than all other constructs (29 ng/ml vs. 0.3 ng/ml for FVIII) (see Figure 7).

[0037] Subject of the invention are, therefore, FVIII-B-domain deleted constructs containing a slightly modified Factor IX truncated intron 1 in different locations of the cDNA. Among these constructs a cDNA bearing the truncated intron 1 in both the FVIII Intron 1 and Intron 12 locations led to a 100 times higher intracellular production than all other constructs and a 9 times higher secretion of the protein. This improved production and secretion was observed in two different cell lines: CHO and HepG2 cells. The FVIII produced was fully active on a one stage clotting assay and appears homogenous on immunoblot. The mRNA amount of all the constructs tested differs no more than three times indicating that the benefit observed in the production is coming both from a transcriptional and a translational effect.

[0038] The present invention indicates that the production of FVIII may be improved by adding introns in the FVIII cDNA. The advantages of such modified FVIII cDNAs for in vitro FVIII production as well as for human gene therapy by inserting such a cDNA in a suitable transvector are important for the future treatment of hemophilia A.

#### Bibliography

- [0039] Dornier, A.J., Bole, D.G., and Kaufman, R.J. (1987): The relationship of N-Linked Glycosylation and Heavy Chain-binding Protein Association with the Secretion of Glycoproteins. *J. Cell, Biol.* **105**, 2665-2674.
- [0040] Fallaux, F.J., Hoeber, R.C., Cramer, S.J., van den Wollenberg, D.J., Briet, E., van Ormondt, H., and van der Eb, A.J. (1996): The human clotting factor VIII cDNA contains an autonomously replicating sequence consensus- and matrix attachment region-like sequence that binds a nuclear factor, represses heterologous gene expression, and mediates the transcriptional effects of sodium butyrate. *Mol Cell Biol* **16**, 4264-72.
- [0041] Foster, P.A., and T.S., Z. (1989): Factor VIII structure and Function. *Blood Reviews* **3**, 180-191.
- [0042] Hoeber, R.C., Fallaux, F.J., Cramer, S.J., van den Wollenberg, D.J., van Ormondt, H., Briet, E., and van der Eb, A.J. (1995): Expression of the blood-clotting factor-VIII cDNA is repressed by a transcriptional silencer located in its coding region. *Blood* **85**, 2447-54.
- [0043] Kaufman, R.J. (1992): Biological regulation of factor VIII activity. *Annu Rev Med* **43**, 325-39.
- [0044] Kaufman, R.J., Wasley, L.C., Davies, M.V., Wise, R.J., Israel, D.I., and Dornier, A.J. (1989): Effects of von Willebrand Factor Coexpression on the Synthesis and Secretion of Factor VIII in Chinese Hamster Ovary Cells. *Mol. Cell. Biol.* **9**, 1233-1242.
- [0045] Koeberl, D.D., Halbert, C.L., Krumm, A., and Miller, A.D. (1995): Sequences within the coding regions of clotting factor VIII and CFTR block transcriptional elongation. *Hum Gene Ther* **6**, 469-79.
- [0046] Kozak, M. (1997): Recognition of AUG and alternative initiator codons is augmented by G in position +4 but is not generally affected by the nucleotides in positions +5 and +6. *EMBO J.* **16**, 2482-92.
- [0047] Kurachi S., Hitomi Y., Furukawa M., and Kurachi K. (1995). Role of the Intron 1 in the expression of the human Factor IX gene. *J. Biol. Chem.* **270**, 5276-5281.
- [0048] Lind, P., Larsson, K., Spira, J., Spira, J., Sysow Backman, M., Almstedt, A., Gray, E., and Sandberg, H. (1995): Novel forms of B-domain-deleted recombinant factor VIII molecules. Construction and biochemical characterization. *Eur J Biochem* **232**, 19-27.
- [0049] Lynch, C.M., Israel, D.I., Kaufman, R.J., and Miller, A.D. (1993): Sequences in the coding region of clotting factor VIII act as dominant inhibitors of RNA accumulation and protein production. *Hum Gene Ther* **4**, 259-72.
- [0050] Nichols, W.C., Seligsohn, U., Zivellin, A., Terry, V.H., Hertel, C.E., Wheatley, M.A., Moussali, M.J., Hauri, H.-P., Ciavarella, N., Kaufman, R.J., and Ginsburg, D. (1998): Mutations in the ER-Golgi Intermediate Compartment Protein ERGIC-53 Cause Combined Deficiency of Coagulation Factors V and VIII. *Cell* **93**, 61-70.
- [0051] Pipe, S.W., Morris, J.A., Shah, J., and Kaufman, R.J. (1998): Differential Interaction of Coagulation Factor VIII and Factor V with Protein Chaperones Calnexin and Calreticulin. *J. Biol. Chem.* **273**, 8567-8544.
- [0052] Pittman D.D., Alerman, E.M., Tomkinson, K.N., Wang, J.H., Giles, A.R., and Kaufman, R.J. (1993): Biochemical, immunological, and in vivo functional characterization of B-domain-deleted factor VIII. *Blood* **81**, 2925-35.
- [0053] Pittman, D.D., Marquette, K.A., and Kaufman, R.J. (1994): Role of the B-domain for factor VIII and factor V



expression and function. *Blood* **84**, 4214-25.

[0054] Vlot, A.J., Koppelman, S.J., Bouma, B.N., and Sixma, J.J. (1998): Factor VIII and von Willebrand Factor. *Thromb. Haemo.* **79**, 456-465.

5

## Annex 1

10

## SEQUENCE LISTING

15

<110> Centeon Pharma GmbH

<120> Modified Factor VIII cDNA

20

<130> C9P28EP

25

<160> 9

<170> PatentIn Ver. 2.1

30

<210> 1

<211> 30

35

<212> DNA

<213> Homo sapiens

40

<400> 1

acacccatgg aaatagagct ctccacctgc

30

45

<210> 2

<211> 24

50

<212> DNA

<213> Homo sapiens

55

5                   <400> 2  
                  agtcctgaag ctagatctct ctcc                   24

10                  <210> 3  
                  <211> 26  
                  <212> DNA  
                  <213> Homo sapiens

15                  <400> 3  
                  aatatggaga gagatctagc tcagg                   26

20                  <210> 4  
                  <211> 36  
25                  <212> DNA  
                  <213> Homo sapiens

30                  <400> 4  
                  ttctcgtcga cgtcttcttg gtcaatggc attggt                   36

35                  <210> 5  
                  <211> 34  
40                  <212> DNA  
                  <213> Homo sapiens

45                  <400> 5  
                  aagacgtcga cgagaaataa ctcgtactac tctt                   34

50                  <210> 6  
                  <211> 26  
55                  <212> DNA

<213> Homo sapiens

5 <400> 6  
agcatgtaga tgctcgccaa taaggc 26

10 <210> 7  
<211> 28

15 <212> DNA  
<213> Homo sapiens

20 <400> 7  
atttggcggg tggaatgcct tattggcg 28

25 <210> 8  
<211> 33

30 <212> DNA  
<213> Homo sapiens

35 <400> 8  
acacctcgag tcagtagagg tcctgtgcct cgc 33

40 <210> 9  
<211> 312

45 <212> DNA  
<213> Homo sapiens

50 <400> 9  
gtttgtttat gcatcctttt ttaaaataca ttgagtatgc ttgcctttta gatatagaaa 60  
tatctgatgc tgtctcttc actaaattt gattacatga ttgacagca atattgaaga 120  
55 gtctaacagc cagcacgcag gttggttaagt actgtgggaa catcacagat ttggctcca 180

tgccctaaag agaaattggc ttccagatta ttggattaa aaacaaagac ttcttaaga 240  
gatgtaaaat ttcatgatg ttctcttt tgctaaaact aaagaattaa cgcgtattct 300  
5 ttacatttc ag 312

10 **Claims**

1. Modified Factor VIII cDNA, **characterised in** that the B-domain of the wild-type Factor VIII cDNA has been deleted and a truncated Factor IX intron 1 has been inserted in one or more locations of the Factor VIII cDNA.
- 15 2. Modified Factor VIII cDNA as claimed in claim 1, **characterised in** that the B-domain of the wild-type Factor VIII cDNA has been replaced by four arginines.
3. Modified Factor VIII cDNA as claimed in claims 1 and 2, **characterised in** that the truncated Factor IX intron 1 has been inserted in the Factor VIII introns 1 and/or 11 and/or 12.
- 20 4. Process for the production of Factor VIII, **characterised in** that the production is performed in a cell-line containing a modified Factor VIII cDNA as claimed in claims 1 to 3.
- 25 5. Transfervector for use in the human gene therapy, **characterised in** that it comprises the modified Factor VIII cDNA of claims 1 to 3.

30

35

40

45

50

55

Fig.1

Schematic view of FLX intron 1 inserted in FVIII intron 1 location




Fig.2

Cloning strategy of Factor IX intron 1 in intron 11 and 12 locations




Fig.3

Kinetic of FVIII production from different transfected cell lines




Fig.4

Procoagulant activities from different transfected cell supernatants




Fig.5

FVIII antigen production in HepG2 cell line  
(average of at least three experiments)






Fig.6

Quantitative analysis of FVIII mRNA in transfected CHO  
(average of at least four different analyses)





**Fig.7**  
**Intracellular amount of FVIII in different cell lines**





European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number  
EP 99 10 4050

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION
A,D	KURACHI S ET AL: "ROLE OF INTRON I IN EXPRESSION OF THE HUMAN FACTOR IX GENE" JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 270, no. 10, 10 March 1995 (1995-03-10), pages 5276-5281, XP002055415 ISSN: 0021-9258 * abstract; figure 2 *		C12N15/12 C07K14/755
A	WO 92 16557 A (KABI PHARMACIA AB) 1 October 1992 (1992-10-01) * page 4, paragraph 1; claim 6; figure 1 *		
A,D	LIND, P. ET AL.: "Novel forms of B-domain-deleted recombinant factor VIII molecules. Construction and biochemical characterization." EUROPEAN JOURNAL OF BIOCHEMISTRY, vol. 232, no. 1, August 1995 (1995-08), pages 19-27, XP002111900 * abstract *		
			TECHNICAL FIELDS SEARCHED
			C07K C12N
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 12 August 1999	Examiner Smalt, R
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 99 10 4050

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

12-08-1999

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9216557 A	01-10-1992	SE 468050 B	26-10-1992
		AU 650730 B	30-06-1994
		AU 1351392 A	21-10-1992
		CA 2081659 A	16-09-1992
		EP 0533862 A	31-03-1993
		FI 925131 A	11-11-1992
		JP 2655752 B	24-09-1997
		PT 100245 A	30-07-1993
		SE 9100799 A	16-09-1992
		US 5661008 A	26-08-1997
-----			

56904 004 0045

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

